



# Micro 600™

## Directions For Use General Guideline:

For proper agronomic application rates suitable for your geographical area or the maximum allowable non-nutrient application rate per acre, consult a trained soil specialist at AgroLiquid or call or write to AgroLiquid with the address provided.

This combination of micronutrients, plus the essential secondary nutrient sulfur, guards against a variety of common deficiencies, including boron, which can affect pollination, and iron, which can damage plant leaves, stalks, buds, and fruit. The Nutriq technology contained in Micro 600 allows for improved uptake and assimilation by the crop.

Crop	In-Furrow
Corn (Grain) 30" Row Spacing	0-1 gal/A
Corn (Silage) 30" Row Spacing	0-1 gal/A
Soybeans 15" Row Spacing	0-0.5 gal/A
Soybeans 30" Row Spacing	0-0.5 gal/A
Sorghum	0-0.5 gal/A
Dry Beans	0-0.25* gal/A
Cotton	0-0.25* gal/A
Sugarbeet	0-0.25* gal/A
Canola	0-0.25* gal/A
Wheat (Spring or Winter)	0-0.25* gal/A
Potato	0-1 gal/A Direct contact with the seed piece
Alfalfa	0-0.25* gal/A

### Foliar Application Recommendations

**RATE: 0.125-2 gallon/acre**

unless otherwise noted

<b>Corn</b> Sidedress	<b>Apples</b> Banded or through drip irrigation during the growing season
<b>Sorghum</b> Sidedress	<b>Tobacco</b> Banded or through drip irrigation during the growing season
<b>Cotton</b> Sidedress	<b>Tree Nuts</b> Banded or through drip irrigation during the growing season;
<b>Sugarbeet</b> Sidedress	<b>Other Tree Fruits</b> Banded or through drip irrigation during the growing season
<b>Wheat</b> Topdress up to Feekes Stage 4	<b>Vegetables</b> Broadcast, surface banded or through drip irrigation during the growing season
<b>Potato</b> Sidedress or fertigation	
<b>Alfalfa</b> Prior to, or within 14 days of spring green-up, and/or 0-7 days after cutting, broadcast	
<b>Grapes</b> Broadcast, surface banded or through drip irrigation during the growing season	
<b>Tomato</b> Banded or through drip irrigation during the growing season	

### Foliar Application Recommendations

**RATE: 0.125-.05 gallon/acre**

unless otherwise noted

<b>Corn</b> <b>Soybean 30" and 15" Rows</b>
<b>Sorghum</b>
<b>Dry Beans</b>
<b>Cotton</b>
<b>Sugarbeet</b>
<b>Canola</b>
<b>Wheat</b>
<b>Potato</b>
<b>Alfalfa</b>
<b>Grapes</b>
<b>Tomato</b>
<b>Tobacco</b>
<b>Apples</b>
<b>Tree Nuts</b>
<b>Other Tree Fruits</b>
<b>Vegetables</b>

Broadcast, or banded not less than 2" from the seed furrow, surface banded, or applied through drip irrigation at the base of the plant

**RATE: 0.125-2 gallon/acre**

Corn	Canola	Tobacco
Soybean	Wheat	Apples
Sorghum	Potato	Tree Nuts
Dry Beans	Alfalfa	Tree Fruit
Cotton	Grapes	Vegetables
Sugarbeet	Tomato	

**0.25% in Transplant Solution**

Grapes	Apples	Vegetables
Tomato	Tree Nuts	
Tobacco	Tree Fruit	

Please consult with an AgroLiquid Sales Account Manager or Agronomist for further direction when utilizing rates higher than the lower limit of the given range.

 **Guaranteed by: AgroLiquid**  
 Division of COG Marketers, Ltd.  
 3055 W M-21  
 St. Johns, MI 48879  
 AGROLIQUID [agroliquid.com](http://agroliquid.com)

NOTE: Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm>



**GS07**  
**WARNING - Irritant: Eyes, Skin**